

Type 2 diabetes-related variants influence the risk of developing multiple myeloma: results from the IMMEnSE consortium

Rafael Ríos^{1,2,*}, Carmen Belén Lupiáñez^{1,2,*}, Daniele Campa^{3,4}, Alessandro Martino³, Joaquin Martínez-López⁵, Manuel Martínez-Bueno⁶, Judit Varkonyi⁷, Ramón García-Sanz⁸, Krzysztof Jamrozik⁹, Charles Dumontet¹⁰, Andrés Jerez Cayuela¹¹, Marzena Wętek¹², Stephano Landi⁴, Anna Maria Rossi⁴, Fabienne Lesueur¹³, Rui Manuel Reis^{14,15,16}, Victor Moreno¹⁷, Herlander Marques^{14,15}, Artur Jurczynski¹⁸, Vibeke Andersen^{19,20}, Ulla Vogel²⁰, Gabriele Buda²¹, Enrico Orciuolo²¹, Svend E H Jacobsen²², Mario Petrini²¹, Annette J Vangsted²³, Federica Gemignani⁴, Federico Canzian³, Manuel Jurado^{1,2} and Juan Sainz^{1,2}

¹Genomic Oncology Area, GENYO, Centre for Genomics and Oncological Research: Pfizer/ University of Granada/Andalusian Regional Government, PTS, Granada, 18016 Granada, Spain

²Hematology Department, Virgen de las Nieves University Hospital, Granada, Spain

³Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany

⁴Department of Biology, University of Pisa, Pisa, Italy

⁵Department of Hematology, Hospital Universitario Doce de Octubre, Madrid, Spain

⁶Area of Genomic Medicine, GENYO, Centre for Genomics and Oncological Research: Pfizer/ University of Granada/Andalusian Regional Government, PTS Granada, Granada, Spain

⁷Semmelweis University, Budapest, Hungary

⁸Haematology Department, University Hospital of Salamanca and IBSAL, Salamanca, Spain

⁹Medical University of Lodz, Lodz, Poland

¹⁰INSERM UMR 1052/CNRS 5286, Université Claude Bernard Lyon I, Lyon, France

¹¹Morales Meseguer General University Hospital, Murcia, Spain

¹²Haematology Clinic, Holly Cross Cancer Center, Kielce, Poland

¹³INSERM, U900, Genetic Epidemiology of Cancers team, Institut Curie, Mines ParisTech, Paris, France

¹⁴Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Braga, Portugal

¹⁵ICVS/3B's – PT Government Associate Laboratory, Braga/Guimarães, Portugal

¹⁶Molecular Oncology Research Center, Barretos Cancer Hospital, Barretos, Brazil

¹⁷IDIBELL – Catalan Institute of Oncology, University of Barcelona, Barcelona 08907, Spain

¹⁸Department of Hematology, Cracow University Hospital, Cracow, Poland

¹⁹Organ Center, Hospital of Southern Jutland, DK-6200 Aabenraa, Denmark

²⁰Faculty of Health Sciences, Institute of Regional Health Research, University of Southern Denmark, DK-5000 Odense C, Denmark

²¹UO Hematology, Department of Internal and Experimental Medicine, University of Pisa, Pisa, Italy

²²Clinic of Biochemistry and Immunology, Laboratory Center, Hospital of Southern Jutland, Aabenraa, Denmark

²³Department of Haematology, Rigshospitalet and Roskilde Hospital, Copenhagen University, Copenhagen, Denmark

* (R Ríos and C B Lupiáñez contributed equally to this work)

Correspondence
should be addressed
to J Sainz
Email
juan.sainz@genyo.es

Abstract

Type 2 diabetes (T2D) has been suggested to be a risk factor for multiple myeloma (MM), but the relationship between the two traits is still not well understood. The aims of this study were to evaluate whether 58 genome-wide-association-studies (GWAS)-identified common variants for T2D influence the risk of developing MM and to determine whether predictive models built with these variants might help to predict the disease risk. We conducted a case-control study including 1420 MM patients and 1858 controls ascertained through the International Multiple Myeloma (IMMEnSE) consortium. Subjects carrying the *KCNQ1*_{rs2237892T} allele or the *CDKN2A-2B*_{rs2383208G/G}, *IGF1*_{rs35767T/T} and *MADD*_{rs7944584T/T} genotypes had a significantly increased risk of MM (odds ratio (OR) = 1.32–2.13) whereas those carrying the *KCNJ11*_{rs5215C}, *KCNJ11*_{rs5219T} and *THADA*_{rs7578597C} alleles or the *FTO*_{rs8050136A/A} and *LTA*_{rs1041981C/C} genotypes showed a significantly decreased risk of developing the disease (OR = 0.76–0.85). Interestingly, a prediction model including those T2D-related variants associated with the risk of MM showed a significantly improved discriminatory ability to predict the disease when compared to a model without genetic information (area under the curve (AUC) = 0.645 vs AUC = 0.629; $P = 4.05 \times 10^{-06}$). A gender-stratified analysis also revealed a significant gender effect modification for *ADAM30*_{rs2641348} and *NOTCH2*_{rs10923931} variants ($P_{\text{interaction}} = 0.001$ and 0.0004, respectively). Men carrying the *ADAM30*_{rs2641348C} and *NOTCH2*_{rs10923931T} alleles had a significantly decreased risk of MM whereas an opposite but not significant effect was observed in women (OR_M = 0.71 and OR_M = 0.66 vs OR_W = 1.22 and OR_W = 1.15, respectively). These results suggest that T2D-related variants may influence the risk of developing MM and their genotyping might help to improve MM risk prediction models.

Key Words

- multiple myeloma
- diabetes
- genetic variants
- susceptibility

Endocrine-Related Cancer
(2015) 22, 545–559

Introduction

Multiple myeloma (MM) is a plasma-cell neoplasm of complex aetiology that may arise as a result of the interaction between adverse environmental and inherited genetic risk factors (Morgan *et al.* 2012). Although survival rates for MM have improved dramatically during the last two decades, likely due to the introduction of novel targeted therapies (proteasome inhibitors, immunomodulators and others), the disease outcome still remains poor with a 5-year overall survival rate not higher than 55% (Kumar *et al.* 2014).

Age, male gender, African ancestry and monoclonal gammopathy of uncertain significance (MGUS) have been established as major risk factors for MM (Alexander *et al.* 2007). In addition, exposure to a wide range of toxins as well as type 2 diabetes (T2D) and obesity have been suggested as important mediators of the complex process of myelomagenesis (Alexander *et al.* 2007, Lope *et al.* 2008, Wallin & Larsson 2011). Among these latter preventable factors, T2D has attracted significant attention since it has been consistently identified as a medical condition frequently found in MM patients (Khan *et al.* 2008, Richardson *et al.* 2009, Castillo *et al.* 2012) and it is thought to influence the myelomagenesis through hyperglycaemia

and insulin-dependent and -independent mechanisms (Xu *et al.* 2014). In a recent well-powered meta-analysis, Castillo *et al.* (2012) observed that T2D was significantly associated with an increased risk of developing the disease (Castillo *et al.* 2012). This finding concurs with those previously reported in several epidemiological studies that showed a high incidence of T2D among MM patients ranging between 11 and 22% (Richardson *et al.* 2006, Badros *et al.* 2007). In addition, it has been reported that T2D may have a negative impact on MM prognosis (Chiu *et al.* 2006, Wu *et al.* 2014) and that the treatment with anti-diabetic drugs may effectively kill MM cells (Wu *et al.* 2014).

Considering that T2D and MM have strong genetic components and share several biological pathways and markers (Xu *et al.* 2014) and that T2D-related polymorphisms may influence the risk of developing solid cancer (Folsom *et al.* 2008, Cheng *et al.* 2011, Sainz *et al.* 2012, Ma *et al.* 2014), we hypothesized that genetic risk factors for T2D may be associated with the risk of developing MM. So far there have not been studies evaluating the impact of diabetogenic variants on the risk of developing hematological cancers. Therefore, we decided to conduct a

multi-centre case-control study including 1420 MM patients and 1858 controls to evaluate whether 58 variants convincingly shown to be associated with T2D contribute to the risk of developing MM. We also aimed at determining whether predictive models including T2D-related variants significantly improve the discriminatory ability to predict the risk of MM.

Material and methods

Study population

The study population consisted of 1420 MM patients (705 women and 715 men) and 1858 controls (916 women and 942 men) ascertained through the International Multiple Myeloma (IMMEnSE) consortium (Supplementary Table 1, see section on supplementary data given at the end of this article), which has been described in detail elsewhere (Martino *et al.* 2012). The diagnosis of MM was assigned by physician and fulfilled the International Myeloma Working Group (IMWG) criteria (International Myeloma Working Group 2003). Controls were blood donors or hospitalized subjects with a diagnosis not related to cancer who were recruited in the same geographical area of the cases (Supplementary Table 1). Additional information concerning to the recruitment strategy of controls is shown in the Supplementary Material. The investigation was approved by the ethical committee of each participant institution, functioning according to the third edition of the Guidelines on the Practice of Ethical Committees in Medical Research issued by the Royal College of Physicians of London (www.rcplondon.ac.uk) and all participants gave their written informed consent to participate in the study.

SNP selection and genotyping

Fifty-eight genome-wide-association-studies (GWAS)-identified variants for T2D were selected to be genotyped in the IMMEnSE consortium population (Table 1 and Supplementary Material). The genotyping of the selected polymorphisms was carried out at GENYO (Centre for Genomics and Oncological Research: Pfizer/University of Granada/Andalusian Regional Government, Granada, Spain) using KASPar assays (LGC Genomics, Hoddesdon, UK) according to manufacturer's instructions. For internal quality control, 5% of samples were randomly selected and included as duplicates. Concordance between the original and the duplicate samples for the 58 SNPs was $\geq 99.0\%$. Call rates for all SNPs were $\geq 90.0\%$ with the exception of the *WFS1*_{rs734312} SNP that was excluded from further analyses.

Statistical analysis

The Hardy-Weinberg Equilibrium (HWE) tests were performed in the control group by a standard observed-expected χ^2 test. Logistic regression analyses were used to assess the effects of the genetic polymorphisms on MM risk using co-dominant, dominant, recessive and log-additive inheritance models. Overall analyses were adjusted for age at diagnosis, gender and country of origin. All analyses were conducted using the statistical software SSPS (version 20.0). Statistical power was calculated using the Quanto vs12.4 (<http://biostats.usc.edu/software>) assuming a log-additive model.

In order to account for multiple testing, we calculated an adjusted significance level using the M_{eff} method (Nyholt 2004), which considers the number of independent marker loci ($M_{\text{eff},i}=55$) but also the number of models of inheritance tested (co-dominant, dominant, recessive and log-additive). Thus, the resulting threshold for the main effect analysis was 0.00022 ($(0.05/55)/4$) (Supplementary Material). Since a study-wide significance threshold considering all these factors is generally perceived as a 'too conservative' test, we also assessed the magnitude of observed associations between selected SNPs and risk of MM through a quantile-quantile (QQ) plot generated from the results of the IMMEnSE population. The observed association P values were ranked in order from smallest to largest on the y -axis and plotted against the expected results from a theoretical $\sim\chi^2$ -distribution under the null hypothesis of no association on the x -axis. A deviation from the identity line would confirm that the number of corresponding associations is more than expected under the null hypothesis and therefore that these associations are likely to be true associations.

Predictive models and discriminative accuracy

We also examined the value of T2D-related polymorphisms for prediction of MM using stepwise logistic and Cox regression analyses. We built a prediction model including age and sex and those genetic variants that showed significant associations with MM in the single-SNP analysis ($P<0.05$). Then, using P values as a selection criterion, we dropped variables that have the highest P value and we stopped when all variables were significant defined by $P<0.10$. The area under the curve (AUC) of a receiver operating characteristic (ROC) curve analysis was used to assess the discriminative accuracy of this particular model compared with a reference model including age and sex as covariates. A $-2\log$ likelihood ratio (LR) test

Table 1 Selected type-2 diabetes-related SNPs

Gene name	dbSNP rs#	Nucleotide substitution	Reference allele IMMEnSE	GWAS- identified risk allele for T2D	Location/Aa substitution	References
ADAM30	rs2641348	T/C ^a	T	C	L359P	Zeggini et al. (2008) and Lyssenko et al. (2009)
ADAMTS9	rs4607103	T/C	C	C	Near gene	Mohlke et al. (2008), Zeggini et al. (2008) and Shu et al. (2010)
ADCY5	rs11708067	T/C	T	T	Intronic	Dupuis et al. (2010) and Saxena et al. (2010)
ADRA2A	rs10885122	G/T	G	G	Near ADRA2A	Dupuis et al. (2010)
ARAP1, CENTD2	rs1552224	G/T	T	A	Near gene	Voight et al. (2010) and Nielsen et al. (2011)
BCL11A	rs10490072	C/T	T	T	Near gene	Zeggini et al. (2008)
CDC123	rs12779790	A/G	A	G	Near gene	Mohlke et al. (2008), Zeggini et al. (2008) and Shu et al. (2010)
CDKAL1	rs7754840	C/G	G	C	Intronic	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Florez et al. (2007) and Scott et al. (2007)
CDKN2A-2B	rs564398	T/C	T	T	Near gene	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Scott et al. (2007), Mohlke et al. (2008), Zeggini et al. (2008), Takeuchi et al. (2009), Shu et al. (2010) and Yamauchi et al. (2010)
CDKN2A-2B	rs10811661	T/C	T	T	Near gene	
CDKN2A-2B	rs2383208	A/G ^b	A	A	Near gene	
COL5A1	rs4240702	C/T	C	NS	Intronic	Bouatia-Naji et al. (2009)
CRY2	rs11605924	A/C	C	A	Intronic	Dupuis et al. (2010)
DCD	rs1153188	A/T	A	A	Near gene	Zeggini et al. (2008)
EXT2	rs1113132	C/G	C	C	Intronic	Florez et al. (2007) and Sladek et al. (2007)
FADS1	rs174550	C/T	A	T	Intronic	Dupuis et al. (2010)
FAM148B	rs11071657	A/G	A	A	Near gene	Chambers et al. (2008) and Dupuis et al. (2010)
FLJ39370	rs17044137	A/T	T	A	Near gene	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007)
FTO	rs8050136	A/C ^c	C	A	Intronic	Wellcome Trust Case Control (2007), Zeggini et al. (2007) and Mohlke et al. (2008)
G6PC2	rs560887	G/A	G	G	Intronic	Bouatia-Naji et al. (2008, 2009), Chen et al. (2008), Prokopenko et al. (2009) and Dupuis et al. (2010)
GCK	rs1799884	G/A	G	A	Near gene	Bouatia-Naji et al. (2008, 2009), Chen et al. (2008), Prokopenko et al. (2009) and Dupuis et al. (2010)
GCKR	rs1260326	C/T	C	T	L445P	Bouatia-Naji et al. (2009), Dupuis et al. (2010) and Saxena et al. (2010)
HHEX	rs1111875	G/A	C	C	Near gene	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Florez et al. (2007), Scott et al. (2007), Sladek et al. (2007), Wellcome Trust Case Control (2007), Zeggini et al. (2007) and Mohlke et al. (2008)
HMGA2	rs1531343	C/G	G	C	Near gene	Voight et al. (2010) and Nielsen et al. (2011)

Table 1 Continued

Gene name	dbSNP rs#	Nucleotide substitution	Reference allele IMMEnSE	GWAS-identified risk allele for T2D	Location/Aa substitution	References
<i>HNF1A, TCF1</i>	rs7957197	A/T	T	T	Intronic	Voight et al. (2010) and Nielsen et al. (2011)
<i>IGF1</i>	rs35767	C/T ^d	C	C	Near gene	Pechlivanis et al. (2007) and Dupuis et al. (2010)
<i>IGF2BP2</i>	rs4402960	G/T	C	T	Intronic	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Florez et al. (2007), Scott et al. (2007), Wellcome Trust Case Control (2007), Zeggini et al. (2007), Mohlke et al. (2008) and Shu et al. (2010)
<i>IL13</i>	rs20541	C/T	C	T	R144Q	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007)
<i>IRS1</i>	rs2943641	C/T	C	C	Near gene	Rung et al. (2009), Voight et al. (2010) and Tang et al. (2013)
<i>JAZF1</i>	rs864745	A/G	A	T	Intronic	Zeggini et al. (2008) and Shu et al. (2010)
<i>KCNJ11</i>	rs5215	T/C ^e	T	C	V337I	Gloyn et al. (2003), Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Scott et al. (2007), Wellcome Trust Case Control (2007), Zeggini et al. (2007), Willer et al. (2007) and Mohlke et al. (2008)
<i>KCNJ11</i>	rs5219	C/T ^f	C	T	K23E	
<i>KCNQ1</i>	rs2237897	C/T	T	C	Intronic	Unoki et al. (2008), Yasuda et al. (2008), Tsai et al. (2010) and Yamauchi et al. (2010)
<i>KCNQ1</i>	rs2074196	G/T	G	G	Intronic	
<i>KCNQ1</i>	rs2237892	C/T ^g	C	C	Intronic	
<i>KCNQ1</i>	rs2237895	A/C	A	C	Intronic	
<i>KCNQ1OT1</i>	rs231362	C/T	C	G	Intronic	Tsai et al. (2010), Voight et al. (2010) and Nielsen et al. (2011)
<i>LTA</i>	rs1041981	A/C ^e	A	A	T60N	Hamid et al. (2005)
<i>MADD</i>	rs7944584	A/T ^d	A	A	Intronic	Dupuis et al. (2010)
<i>MCR4</i>	rs12970134	A/G	G	A	Near gene	Chambers et al. (2008)
<i>MTNR1B</i>	rs1387153	C/T	C	T	Near gene	Bouatia-Naji et al. (2009), Prokopenko et al. (2009) and Voight et al. (2010)
<i>NOTCH2</i>	rs10923931	G/T ^h	G	T	Intronic	Mohlke et al. (2008) and Zeggini et al. (2008)
<i>PKN2</i>	rs6698181	C/T	C	T	Intergenic	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007)
<i>PPARG</i>	rs1801282	C/G	C	C	P12A	Altshuler et al. (2000), Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Scott et al. (2007), Wellcome Trust Case Control (2007), Willer et al. (2007), Zeggini et al. (2007, 2008) and Mohlke et al. (2008)
<i>PRC1</i>	rs8042680	A/C	C	A	Intronic	Voight et al. (2010) and Nielsen et al. (2011)
<i>PROX1</i>	rs340874	A/G	A	G	Promoter	Dupuis et al. (2010)
<i>RBMS1</i>	rs7593730	C/T	C	T	Intronic	Qi et al. (2010)
<i>SLC2A2</i>	rs11920090	A/T	A	T	Intronic	Dupuis et al. (2010)

Table 1 Continued

Gene name	dbSNP rs#	Nucleotide substitution	Reference allele IMMEnSE	GWAS- identified risk allele for T2D	Location/Aa substitution	References
<i>SLC30A8</i>	rs13266634	C/T	C	C	R325W	Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Florez et al. (2007), Scott et al. (2007), Sladek et al. (2007), Steinthorsdottir et al. (2007), Wellcome Trust Case Control (2007), Zeggini et al. (2007), Mohlke et al. (2008), Dupuis et al. (2010) and Shu et al. (2010)
<i>TCF2</i>	rs7501939	C/T	C	C	Intronic	Gudmundsson et al. (2007) and Sandhu et al. (2007)
<i>TCF7L2</i>	rs7903146	C/T	C	T	Intronic	Grant et al. (2006), Scott et al. (2006), Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research et al. (2007), Florez et al. (2007), Scott et al. (2007), Sladek et al. (2007), Steinthorsdottir et al. (2007), Wellcome Trust Case Control (2007), Zeggini et al. (2007), Mohlke et al. (2008), Dupuis et al. (2010) and Saxena et al. (2010)
<i>TCF7L2</i>	rs12255372	G/T	G	T	Intronic	
<i>THADA</i>	rs7578597	T/C ^e	T	T	T1187A	Zeggini et al. (2008)
<i>TP53INP1</i>	rs896854	A/G	G	G	Intronic	Voight et al. (2010) and Nielsen et al. (2011)
<i>TSPAN8</i>	rs7961581	C/T	T	C	Near gene	Grarup et al. (2008)
<i>VEGFA</i>	rs9472138	C/T	C	T	Near gene	Zeggini et al. (2008)
<i>WFS1</i>	rs734312	A/G	A	NS	H611R	Sandhu et al. (2007)
<i>WFS1</i>	rs10010131	A/G	G	G	Intronic	Sandhu et al. (2007)

NS, not specified; Aa, aminoacid; GWAS, genome-wide association studies. Effect allele in bold and underlined.

^aC allele was associated with a decreased risk of MM in men whereas an opposite effect was detected in women.

^bG allele was associated with an increased risk of developing MM.

^cA/A genotype was associated with a decreased risk of MM (recessive model).

^dT/T genotype was associated with a decreased risk of MM (recessive model).

^eC allele was associated with a decreased risk of MM.

^fT allele was associated with an increased risk of MM.

^gT allele was associated with an increased risk of MM.

^hT allele was associated with a decreased risk of MM in men whereas an opposite effect was detected in women.

was used to determine whether the predictive model including genetic information fitted significantly better the data compared to the reference model. Although the addition of genetic variables to the reference model will almost always make the model fit better, the LR test allowed us to confirm whether the difference in model fit between both models was statistically significant. Besides this suggestive analysis, we also ran a randomization test to confirm whether the improved predictive ability of the model including genetic variants significantly associated with MM was consistent after 10 000 iterations. We compared our full predictive model including significant SNPs, age and gender ('original' model) with 10 000 'randomized' models in which the effect of SNPs on MM risk was neutralized by reassigning randomly all genotypes

(null distribution; [Supplementary Material](#)). Subsequently, we calculated an empirical $P_{\text{iterations}}$ -value by dividing the number of times in which the 'randomized' AUC value was equal or greater than the 'original' AUC value by the number of iterations. Then, we could also calculate the Z score and $P_{\text{Z score}}$ -value for the original AUC using the 'randomized' AUC average of these 10 000 iterations and their s.d. All analyses were performed using R software (<http://www.r-project.org/>).

Gender-specific association analysis

We also evaluate gender-specific associations of selected SNPs with MM risk. Logistic regression analyses were corrected for age and country of origin. Of note, to

evaluate whether a different gender distribution across populations within the IMMEnSE consortium could be responsible for the gender effect modification observed for certain SNPs, we also assessed heterogeneity and index I^2 statistic using Cochran's χ^2 based Q statistic test (Lau et al. 1997). Heterogeneity was considered significant when $P < 0.1$.

Results

Overall associations of selected SNPs with MM risk

All SNPs were in HWE ($P > 0.001$) with the exception of the *COL5A1*_{rs4240702}, which was therefore excluded from the statistical analyses. Logistic regression analysis showed that carriers of the *KCNQ1*_{rs2237892T} allele or the *CDKN2A-2B*_{rs2383208G/G}, *IGF1*_{rs35767T/T} and *MADD*_{rs7944584T/T} genotypes had an increased risk of MM (odds ratio (OR)=1.32, 95% CI 1.01–1.71, $P=0.039$; OR=1.86, 95% CI 1.12–3.11, $P=0.016$; OR=2.13, 95% CI 1.35–3.37, $P=0.0012$ and OR=1.33, 95% CI 1.06–1.67, $P=0.014$, respectively) whereas those harbouring the *KCNJ11*_{rs5215C}, *KCNJ11*_{rs5219T} and *THADA*_{rs7578597C} alleles or the *FTO*_{rs8050136A/A} and *LTA*_{rs1041981C/C} genotypes showed a decreased risk for the disease (OR=0.85, 95% CI 0.73–0.99, $P=0.038$; OR=0.84, 95% CI 0.72–0.99, $P=0.034$; OR=0.81, 95% CI 0.68–0.98, $P=0.032$; OR=0.78, 95% CI 0.64–0.95, $P=0.013$ and OR=0.76, 95% CI 0.58–0.99, $P=0.042$, respectively; Table 2). When we corrected for multiple testing (with a threshold of $P=0.00022$), none of the reported associations remained statistically significant. The strongest association observed was for the *IGF1*_{rs35767} SNP with an increased risk of developing MM (OR=2.13, 95% CI 1.35–3.37, $P=0.0012$). In spite of these results, the QQ plot showed an early deviation of identity line, which suggested a high proportion of true associations for a given P value (Fig. 1). Therefore, the data suggest that the effect attributed to SNPs in T2D-related loci (*FTO*, *MADD*, *CDKN2A-2B*, *LTA*) might represent true associations.

Predictive value of T2D-related variants

In order to determine whether there was a joint effect of SNPs significantly associated with MM, we built a prediction model including gender and those nine SNPs showing overall significant associations with MM. After excluding the variables that did not remain significant in the model, the final model included six SNPs that increased the discriminatory ability to predict the risk of

MM when compared with a reference model including age and gender as covariates (AUC=0.645 95% CI 0.624–0.666; Table 3). The LR test showed that the model including genetic variants fitted better the data than the reference model, and that the difference in model fit between both models was statistically significant ($P=4.05 \times 10^{-06}$). In addition, when we evaluated whether the model including genetic variants was consistent in predicting better the MM risk, we found that it showed an AUC value systematically higher than those of the 10 000 randomized models (null distribution; Z score=6.42, $P=6.81 \times 10^{-11}$; Supplementary Material), emphasizing the importance of considering genetic variants significantly associated with MM when building predictive models.

Gender-specific associations with MM risk

Interestingly, a gender-stratified analysis also revealed significant gender effect modifications for *ADAM30*_{rs2641348} and *NOTCH2*_{rs10923931} SNPs ($P_{\text{interaction}}=0.001$ and 0.0004 respectively). For *ADAM30*_{rs2641348C} and *NOTCH2*_{rs10923931T} alleles, a significantly reduced risk for the disease was observed in men (per-allele OR=0.71, 95% CI 0.54–0.94, $P=0.015$ and per-allele OR=0.66, 95% CI 0.50–0.86, $P=0.0019$, respectively) whereas a non-significant opposite effect was seen in women (per-allele OR=1.22, 95% CI 0.93–1.60 and per-allele OR=1.15, 95% CI 0.89–1.50 respectively). A statistically significant heterogeneity, considering $P < 0.05$ as a threshold, was also confirmed for these two SNPs ($P_{\text{HET}}=0.0039$ and $I^2=87.99\%$ and $P_{\text{HET}}=0.0024$ and $I^2=89.12\%$, respectively), which supports the notion suggesting a role of gender in modulating the effect of these SNPs on MM risk. Although there was not a significant interaction with gender, we observed additional gender-specific associations for *WFS1*_{rs10010131}, *THADA*_{rs7578597}, *EXT2*_{rs1113132} and *GCK*_{rs1799884} SNPs according to dominant or recessive models of inheritance (Table 2 and Supplementary Material).

When we took account of multiple testing (with a threshold of $P=0.00022$), we found that the effect of the *IGF1*_{rs35767} variant was stronger in women than men with an association approaching significance with an increased risk of developing MM (OR=3.13, 95% CI 1.46–6.71, $P=0.0026$ vs OR=1.69, 95% CI 0.94–3.02, $P=0.079$ respectively). In addition, we found that the association of *NOTCH2*_{rs10923931} SNP with a decreased risk of MM in men was close to significance according to dominant and log-additive models (OR=0.66, 95% CI 0.50–0.86,

Table 2 Association of T2D-related variants and risk of developing MM

Variant_dbSNP	Gene	Overall (n = 3278)		Men (n = 1657)		Women (n = 1621)		P _{interaction}
		OR (95% CI) ^a	P value	OR (95% CI) ^b	P value	OR (95% CI) ^b	P value	
rs2641348 ^{cd}	ADAM30	0.94 (0.78–1.14)	0.53	0.71 (0.54–0.94)	0.015	1.22 (0.93–1.60)	0.15	0.001
rs4607103	ADAMTS9	1.00 (0.86–1.16)	1.00	0.97 (0.79–1.20)	0.80	1.05 (0.84–1.30)	0.69	0.803
rs11708067	ADCY5	1.02 (0.88–1.20)	0.77	1.05 (0.85–1.31)	0.64	0.97 (0.77–1.22)	0.79	0.425
rs10885122	ADRA2A	1.08 (0.91–1.28)	0.40	1.16 (0.92–1.47)	0.21	0.99 (0.77–1.28)	0.96	0.494
rs1552224 ^e	ARAP1, CENTD2	1.16 (0.72–1.88)	0.54	1.78 (0.96–3.29)	0.066	0.58 (0.25–1.37)	0.20	0.090
rs10490072	BCL11A	1.01 (0.87–1.17)	0.92	1.01 (0.82–1.25)	0.91	1.01 (0.81–1.25)	0.94	0.691
rs12779790	CDIC123, CAMK1D	0.87 (0.74–1.02)	0.075	0.82 (0.66–1.03)	0.083	0.90 (0.71–1.13)	0.35	0.758
rs7754840	CDKAL1	0.98 (0.85–1.14)	0.84	1.05 (0.86–1.29)	0.63	0.91 (0.74–1.13)	0.40	0.376
rs564398	CDKN2A-2B	0.94 (0.80–1.10)	0.42	0.90 (0.72–1.11)	0.33	0.97 (0.78–1.22)	0.80	0.893
rs10811661	CDKN2A-2B	1.02 (0.87–1.20)	0.79	1.11 (0.89–1.38)	0.35	0.93 (0.74–1.17)	0.52	0.358
rs2383208 ^e	CDKN2A-2B	1.86 (1.12–3.11)	0.016	1.92 (1.03–3.58)	0.039	1.68 (0.69–4.10)	0.25	0.585
rs11605924	CRY2	0.93 (0.79–1.10)	0.40	0.95 (0.75–1.19)	0.64	0.92 (0.72–1.17)	0.49	0.747
rs1153188	DCD	0.91 (0.79–1.06)	0.24	0.83 (0.67–1.02)	0.082	1.05 (0.84–1.30)	0.69	0.072
rs1113132 ^e	EXT2	0.92 (0.68–1.24)	0.57	1.24 (0.83–1.87)	0.30	0.64 (0.41–1.00)	0.046	0.067
rs174550	FADS1	1.11 (0.95–1.28)	0.18	1.13 (0.92–1.38)	0.25	1.08 (0.87–1.34)	0.48	0.359
rs11071657	FAM148B	1.03 (0.88–1.20)	0.73	0.95 (0.77–1.17)	0.62	1.16 (0.93–1.46)	0.18	0.275
rs17044137	FLJ39370	0.91 (0.78–1.05)	0.19	1.05 (0.85–1.29)	0.65	0.89 (0.72–1.11)	0.30	0.357
rs8050136 ^e	FTO	0.78 (0.64–0.95)	0.013	0.70 (0.53–0.93)	0.013	0.88 (0.66–1.17)	0.37	0.420
rs560887 ^e	G6PC2	1.16 (0.88–1.52)	0.30	0.98 (0.67–1.44)	0.93	1.46 (0.98–2.18)	0.065	0.386
rs1799884 ^c	GCK	1.10 (0.93–1.30)	0.29	1.27 (1.01–1.61)	0.044	0.92 (0.72–1.18)	0.51	0.254
rs1260326	GCKR	0.92 (0.78–1.08)	0.32	0.93 (0.74–1.17)	0.53	0.90 (0.71–1.13)	0.36	0.926
rs1111875	HHEX	1.14 (0.98–1.33)	0.093	1.09 (0.88–1.36)	0.41	1.21 (0.97–1.52)	0.095	0.452
rs35767 ^e	IGF1	2.13 (1.35–3.37)	0.0012	1.69 (0.94–3.02)	0.079	3.13 (1.46–6.71)	0.0026	0.538
rs4402960	IGF2BP2	1.05 (0.91–1.22)	0.52	0.95 (0.77–1.16)	0.60	1.16 (0.93–1.44)	0.18	0.304
rs20541	IL13	1.01 (0.86–1.18)	0.91	1.14 (0.92–1.42)	0.22	0.88 (0.70–1.10)	0.26	0.147
rs2943641	IRS1	1.08 (0.93–1.26)	0.31	1.16 (0.94–1.43)	0.17	1.00 (0.80–1.24)	0.99	0.492
rs864745 ^e	JAZF1	0.88 (0.74–1.04)	0.14	0.98 (0.77–1.25)	0.85	0.79 (0.61–1.01)	0.060	0.225
rs5215	KCNJ11	0.85 (0.73–0.99)	0.038	0.89 (0.72–1.10)	0.28	0.82 (0.66–1.02)	0.074	0.795
rs5219	KCNJ11	0.84 (0.72–0.99)	0.034	0.92 (0.74–1.14)	0.43	0.78 (0.62–0.98)	0.033	0.587
rs2237897 ^c	KCNQ1	1.25 (0.97–1.61)	0.081	1.08 (0.74–1.56)	0.69	1.42 (1.00–2.01)	0.052	0.580
rs2074196	KCNQ1	1.01 (0.75–1.37)	0.93	0.80 (0.50–1.27)	0.33	1.21 (0.81–1.81)	0.35	0.456
rs2237892	KCNQ1	1.32 (1.01–1.71)	0.039	1.15 (0.78–1.69)	0.48	1.47 (1.03–2.10)	0.036	0.741
rs2237895	KCNQ1	0.91 (0.77–1.08)	0.28	0.94 (0.75–1.18)	0.59	0.91 (0.71–1.16)	0.44	0.878
rs231362	KCNQ1OT1	1.03 (0.87–1.22)	0.71	1.07 (0.85–1.34)	0.59	1.01 (0.80–1.29)	0.92	0.873
rs1041981 ^e	LTA	0.76 (0.58–0.99)	0.042	0.85 (0.59–1.21)	0.36	0.68 (0.45–1.01)	0.050	0.453
rs7944584 ^e	MADD	1.33 (1.06–1.67)	0.014	1.47 (1.08–2.00)	0.015	1.16 (0.83–1.62)	0.39	0.245
rs12970134	MCR4	0.96 (0.82–1.11)	0.58	1.01 (0.82–1.25)	0.89	0.92 (0.72–1.18)	0.51	0.643
rs1387153	MTNR1B	1.03 (0.89–1.19)	0.73	1.01 (0.82–1.24)	0.91	1.05 (0.85–1.30)	0.67	0.761
rs10923931 ^c	NOTCH2	0.88 (0.73–1.06)	0.16	0.66 (0.50–0.86)	0.0019	1.15 (0.89–1.50)	0.29	0.0004
rs7957197 ^e	HNFA1, OASL	1.33 (0.93–1.92)	0.12	1.60 (0.98–2.60)	0.059	1.04 (0.60–1.81)	0.88	0.388
rs6698181	PKN2	1.00 (0.85–1.17)	0.99	1.10 (0.88–1.36)	0.41	0.90 (0.72–1.12)	0.34	0.450
rs1801282	PPARG	1.06 (0.89–1.26)	0.52	1.02 (0.80–1.30)	0.88	1.11 (0.86–1.43)	0.43	0.655
rs8042680 ^e	PRC1	1.24 (0.99–1.55)	0.056	1.21 (0.89–1.64)	0.24	1.37 (0.99–1.89)	0.055	0.777
rs340874	PROX1	0.95 (0.80–1.13)	0.55	0.90 (0.71–1.14)	0.37	1.02 (0.79–1.30)	0.89	0.387
rs7593730	RBMS1	1.10 (0.95–1.29)	0.20	1.18 (0.95–1.46)	0.13	1.06 (0.85–1.33)	0.58	0.852
rs1531343	RPSAP52, HMGA2	0.96 (0.81–1.15)	0.69	1.07 (0.84–1.37)	0.57	0.86 (0.66–1.11)	0.24	0.288
rs11920090	SLC2A2	1.02 (0.86–1.20)	0.84	1.17 (0.93–1.47)	0.18	0.88 (0.68–1.12)	0.29	0.257

Table 2 Continued

Variant_dbsNP	Gene	Overall (n = 3278)		Men (n = 1657)		Women (n = 1621)	
		OR (95% CI) ^a	P value	OR (95% CI) ^b	P value	OR (95% CI) ^b	P _{interaction}
rs13266634	SLC30A8	0.91 (0.78–1.05)	0.19	0.95 (0.78–1.17)	0.64	0.86 (0.69–1.07)	0.17
rs7501939	TCF2	1.06 (0.91–1.24)	0.43	1.06 (0.85–1.33)	0.58	1.11 (0.89–1.38)	0.37
rs7903146	TCF7L2	0.99 (0.85–1.15)	0.90	1.10 (0.89–1.36)	0.37	0.88 (0.71–1.10)	0.26
rs12255372	TCF7L2	0.94 (0.81–1.09)	0.43	1.06 (0.86–1.30)	0.60	0.83 (0.67–1.03)	0.088
rs7578597 ^c	THADA	0.81 (0.68–0.98)	0.032	0.91 (0.70–1.18)	0.47	0.73 (0.56–0.96)	0.025
rs896854 ^c	TP53INP1	1.17 (0.99–1.39)	0.072	1.22 (0.96–1.55)	0.01	1.13 (0.88–1.45)	0.33
rs7961581	TP53INP1	1.03 (0.89–1.19)	0.71	1.03 (0.84–1.27)	0.78	1.04 (0.84–1.29)	0.71
rs9472138	VEGFA, LGR5	1.13 (0.97–1.31)	0.11	1.19 (0.97–1.46)	0.10	1.09 (0.88–1.35)	0.43
rs734312	WFS1	0.98 (0.83–1.16)	0.84	1.05 (0.83–1.33)	0.67	0.90 (0.71–1.15)	0.39
rs10010131	WFS1	0.94 (0.80–1.10)	0.42	1.11 (0.89–1.38)	0.34	0.77 (0.62–0.96)	0.022

Abbreviations: SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval; NS, not specified. Estimates were adjusted for age, sex, country of origin. $P < 0.05$ in bold.

^aEstimates calculated according to a dominant model of inheritance and adjusted for age, gender and region.

^bEstimates calculated according to a dominant model of inheritance and adjusted for age and region.

^cADAM30_{rs2641348} (per-allele OR_{MEN} = 0.71, 95% CI 0.55–0.92; P_{trend} = 0.0072 vs per-allele OR_{WOMEN} = 1.29, 95% CI 1.00–1.65; P_{trend} = 0.046). GCK_{rs1799884} (per-allele OR_{MEN} = 1.24, 95% CI 1.00–1.54; P_{trend} = 0.050 vs per-allele OR_{WOMEN} = 0.94, 95% CI 0.75–1.18; P_{trend} = 0.58). KCNQ1_{rs2237897} (per-allele OR_{MEN} = 1.09, 95% CI 0.77–1.54; P_{trend} = 0.63 vs per-allele OR_{WOMEN} = 1.41, 95% CI 1.02–1.97; P_{trend} = 0.041). KCNQ1_{rs2237892} (per-allele OR_{MEN} = 1.13, 95% CI 0.77–1.66; P_{trend} = 0.52 vs per-allele OR_{WOMEN} = 1.47, 95% CI 1.04–2.08; P_{trend} = 0.030). NOTCH2_{rs10923831} (per-allele OR_{MEN} = 0.66, 95% CI 0.52–0.84; P_{trend} = 0.0007 vs per-allele OR_{WOMEN} = 1.22, 95% CI 0.96–1.56; P_{trend} = 0.010). THADA_{rs7578597} (per-allele OR_{MEN} = 0.96, 95% CI 0.75–1.22; P_{trend} = 0.73 vs per-allele OR_{WOMEN} = 0.75, 95% CI 0.58–0.96; P_{trend} = 0.021). TP53INP1_{rs896854} (per-allele OR_{MEN} = 1.17, 95% CI 1.01–1.35; P_{trend} = 0.04 vs per-allele OR_{WOMEN} = 1.03, 95% CI 0.89–1.20; P_{trend} = 0.68).

^dADAM30_{rs2641348} (OR_{RECESSIVE-MEN} = 0.36, 95% CI 0.12–1.13; P = 0.060 vs OR_{RECESSIVE-WOMEN} = 4.40, 95% CI 1.44–13.40; P = 0.0059).


^eEstimates calculated according to a recessive model of inheritance.

$P = 0.0019$ and $P_{trend} = 0.0007$), which may suggest a gender-specific allele-dosage effect for this variant to modulate the disease risk ($P_{interaction} = 0.0004$; Table 2). According to a log-additive model, we also found that the association of the ADAM30_{rs2641348C} allele with a decreased risk of MM in men showed a slight trend to be significant considering multiple testing (OR = 0.71, 95% CI 0.55–0.92, $P = 0.0072$) whereas, according to a recessive model, the association of the ADAM30_{rs2641348C/C} genotype was also close to survive multiple testing correction (OR = 4.40, 95% CI 1.44–13.40, $P = 0.00059$; Table 2 and Supplementary Table 2, see section on supplementary data given at the end of this article).

Discussion

In the present study, we report for the first time evidence of significant associations between GWAS-identified T2D genetic variants and MM risk. We found that carriers of the KCNQ1_{rs2237892T} allele, CDKN2A-2B_{rs2383208G/G}, IGF1_{rs35767T/T} and MADD_{rs7944584T/T} genotypes were at increased risk of MM, whereas those carrying the KCNJ11_{rs5215C}, KCNJ11_{rs5219T} and THADA_{rs7578597C} alleles or the FTO_{rs8050136A/A} and LTA_{rs1041981C/C} genotypes showed a decreased risk for the disease. The associations for the KCNQ1, CDKN2A-2B, IGF1, MADD, KCNJ11, and THADA gene variants with the risk of MM showed an opposite direction to those previously reported in the GWAS for T2D (i.e., the risk allele was the opposite for MM and T2DM), which points towards a non-diabetogenic mechanism underlying the effect of these variants to modulate the risk of the disease. In support of this hypothesis, several studies have suggested that, besides their influence on pancreatic function and insulin secretion through a wide variety of biological mechanisms, some of these genes may also act as tumour suppressor genes (Koh *et al.* 1995, Kim & Sharpless 2006, Than *et al.* 2013) and have an impact in the modulation of cell survival (Butt *et al.* 1999, Ortega *et al.* 2002, Sharifi *et al.* 2013), differentiation (Pancewicz *et al.* 2010), proliferation (Grimberg 2003, Pancewicz *et al.* 2010) and apoptosis (LeRoith *et al.* 1995, Li *et al.* 2008, Pancewicz *et al.* 2010). Interestingly, a recent study demonstrated that T2D status was not implicated in the relationship between HNF1B and JAZF1 variants and prostate cancer risk (Stevens *et al.* 2010), which is in line with our hypothesis suggesting that T2D-related variants may determine the risk of MM through non-diabetogenic mechanisms.

When we took into account multiple testing corrections, only the association of the IGF1_{rs35767} promoter

**Figure 1**

QQ plot used to evaluate the magnitude of observed associations of T2D-related variants with risk of MM. QQ plot was calculated assuming a recessive model of inheritance. Deviation from the expected distribution is observed above an expected χ^2 of 0.75. The x-axis is $-\log_{10}$ of the expected P values (under a null hypothesis of no effects) whereas the y-axis is $-\log_{10}$ values of the actual P values.

polymorphism with an increased risk of developing MM remained close to significance ($P=0.0012$), which suggested that the *IGF1* locus may play an important role in triggering cell proliferation in malignant plasma cells. In support of the hypothesis, it has been observed that *IGF1* acts as a major growth factor in MM that, directly or in cooperation with other growth factors, induces MM cell growth and proliferation (Ferlin *et al.* 2000, Bommert *et al.* 2006, Sprynski *et al.* 2009) and can

eventually lead to chemoresistance (Xu *et al.* 1997, Kuhn *et al.* 2012). Likewise, it has been also reported that treatment with metformin, an anti-diabetic drug that inhibits *IGF1* signaling pathway, significantly reduces the risk of transformation from MGUS to symptomatic MM (American Society of Clinical Oncology Annual Meeting 2014; abstract 1532) and that constant use of this treatment may induce cell apoptosis (Rattan *et al.* 2012) and enhance the effectiveness of chemotherapeutic regimes in blood and solid cancers (Feng *et al.* 2011, Pan *et al.* 2012, Watson 2013). Interestingly, several authors have also reported that *IGF1* and its analogues are associated with an increased death in patients with progressive MM (Standal *et al.* 2002, Chou *et al.* 2012, Wu *et al.* 2014), whereas the administration of metformin results in the reduction of deaths in patients with progressive disease (Wu *et al.* 2014).

In fact, Chen *et al.* (2013) recently demonstrated that the *IGF1*_{rs35767} SNP together with two neighbour SNPs constitutes a haplotype that efficiently regulates transcriptional activity (Chen *et al.* 2013). Similarly, several studies have consistently reported that carriers of the *IGF1*_{rs35767T} allele showed significantly higher levels of circulating *IGF1* than those harbouring the WT allele (Mannino *et al.* 2013, Sesti *et al.* 2014) and that the presence of this variant is associated with an increased risk of developing several types of cancer (Ollberding *et al.* 2012, Qian *et al.* 2014).

Although it is tempting to speculate that the *IGF1*_{rs35767} SNP may be responsible for the effect attributed to diabetes on the risk of MM, we believe that rather than

Table 3 Discriminative value AUC for models including T2D-related variants

SNPs	P value	OR 95% CI	AUC 95% CI ^{a,b}
Reference model ^c			
Gender	0.731	0.972 (0.828–1.141)	
Age	$<2.00 \times 10^{-16}$	1.036 (1.030–1.042)	0.629 (0.607–0.650) ^e
Predictive model built with six significant SNPs ^d			
<i>IGF1</i> _{rs35767}	0.004	2.076 (1.258–3.426)	
<i>FTO</i> _{rs8050136}	0.002	0.723 (0.586–0.892)	
<i>MADD</i> _{rs7944584}	0.094	1.218 (0.967–1.535)	
<i>PRC1</i> _{rs8042680}	0.061	1.261 (0.989–1.607)	
<i>KCNJ11</i> _{rs5215}	0.027	0.832 (0.706–0.980)	
<i>KCNQ1</i> _{rs2237892}	0.008	1.468 (1.106–1.950)	
Gender	0.776	1.024 (0.871–1.204)	
Age	$<2.00 \times 10^{-16}$	1.037 (1.031–1.043)	0.645 (0.624–0.666) ^{ef}

^aIncluding age and gender as variables never dropped from models.

^bCompared with a baseline model with AUC=0.5.

^cIncluding age and gender as covariates.

^dSNPs showing a significant association with MM ($P<0.05$). After removing missing values, 2460 subjects were available for prediction capacity analysis.

^eA LR test showed that the model including genetic variants fitted the data better than the reference model and that the difference in model fit between both models was statistically significant ($-2\log$ likelihood ratio test, $df=6$, $P=4.05 \times 10^{-06}$). residual deviance (reference model): 3380.4. residual deviance (significant SNPs model): 3345.2.

^fA sort analysis revealed that this model showed an AUC value systematically higher than those observed in 10 000 randomized models (null distribution; Z score = 6.42, $P=6.81 \times 10^{-11}$; Supplementary Material).

acting separately to modulate the risk of the disease, this genetic variant acts along with additional variants within *KCNQ1*, *CDKN2A-2B*, *MADD*, *KCNJ11*, *THADA*, *LTA* and *FTO* genes to modulate the disease risk. In order to test this hypothesis, we decided to evaluate the predictive value of T2D-related polymorphisms for prediction of MM using stepwise logistic and Cox regression analyses. Interestingly, we found that adding genetic factors to a model without covariates (including only age and gender) substantially improved the prediction of disease development. A predictive model including six SNPs significantly associated with MM in the single analysis showed an adjusted concordance statistic AUC of 64.5% for MM. The consistency of this result was confirmed through a randomization test that showed that none of the 10 000 randomized models showed a higher AUC value than our 'original' model including six genetic variants, age and gender. The addition of genetic variants associated with MM at $P < 0.10$ level did not improve the discriminatory ability to predict MM, which pointed towards a joint contribution of *IGF1*, *FTO*, *MADD*, *PRC1*, *KCNJ11* and *KCNQ1* polymorphisms to predict the risk of the disease. Although the prediction capacity of these models could be considered relatively small when compared with a reference model, the relative absence of current diagnostic factors for MM suggest that the use of genetic variants could be a good option to improve the prediction of the disease risk.

The association of the non-diabetogenic alleles or genotypes for the polymorphisms within *KCNQ1*, *CDKN2A-2B*, *MADD*, *KCNJ11*, *THADA* and *FTO* genes with the risk of MM suggests that these genes, rather than acting through an insulin-dependent mechanism, may modulate the risk of MM by acting as tumour suppressor genes (Duro *et al.* 1995) or through mechanisms promoting cancer cell apoptosis (Rippe *et al.* 2003, Turner *et al.* 2013). In support of this idea, it has been recently reported that most of these genes are highly expressed in tumours and that genetic polymorphisms in these loci are also associated with cancer development (Sauroja *et al.* 2000, Cander *et al.* 2014) and tumour progression (Chen *et al.* 2009).

The association of *LTA*_{rs1041981} SNP with a decreased risk of MM showed a similar direction to that observed in the GWAS for T2D (*LTA*_{rs1041981A} as risk allele) suggests a diabetogenic effect of this SNP to modify the risk of MM. However, we could not dismiss the idea suggesting that observed association could be due to a different distribution of diabetics between MM cases and controls. Similarly, although the direction of the association for the *FTO*_{rs8050136} SNP with the risk of MM was opposite to the one observed in the GWAS for T2D, we could not rule

out the possibility that the observed effect for this obesogenic SNP (Scuteri *et al.* 2007) could be due to significant differences in BMI between MM cases and controls. Further studies using well-characterized cohorts are needed to confirm these latter associations.

Although it was not the primary objective of this study, we also performed gender-stratified analysis to assess whether there was a gender effect modification of selected SNPs to modulate the risk of developing MM. Interestingly, we found a significant gender effect modification for *ADAM30*_{rs2641348}, and *NOTCH2*_{rs10923931} SNPs, which suggested a gender-specific effect of these loci to modulate the risk of MM. We observed that, according to a log-additive model of inheritance, the association of the *NOTCH2*_{rs10923931} SNP with a decreased risk of MM in men and the association of *ADAM30*_{rs2641348C/C} genotype with an increased risk of MM in women showed a marginal level of significance after correction for multiple testing. Recently, it has been demonstrated that *NOTCH2* is highly expressed in MM cells and that it is a key regulator of MM pathogenesis (Colombo *et al.* 2013). In particular, it has been reported that the activation of the *NOTCH2*, which interacts with Wnt components, induces an exacerbated growth of MM cells and accelerated the course of the disease by promoting cancer stem cell self-renewal (Xu *et al.* 2012a) and resistance to chemotherapeutic agents (Xu *et al.* 2012b). Considering that *NOTCH2*_{rs10923931} and *ADAM30*_{rs2641348} are neighbour SNPs in strong linkage disequilibrium (Zeggini *et al.* 2008) and that they showed gender-specific associations with the risk of MM, we hypothesize that *NOTCH2-ADAM30* might represent a gender-specific susceptibility region for MM. In support of this idea, it has been reported that gender-specific variants within *NOTCH* and *WNT* signalling pathways, which are involved in determining cell proliferation and differentiation, may lead to important gender-specific differences in tumour recurrence and chemoresistance (Paez *et al.* 2014).

Our study has both strengths and weaknesses. The major strength is the large sample size. To our knowledge, this is the first study to evaluate the overall and gender-specific associations of T2D-related variants with the risk of developing MM and to assess their predictive value for MM. Although the influence of diabetogenic variants on the risk of the disease was expected to be very modest, our study was sufficiently powered to detect such small effects. Based on the genotype frequencies observed in our study cohort, we had 80% of power (dominant model) to detect an OR of 1.29 at $\alpha = 0.00022$ (multiple testing threshold) for a polymorphism with a minor allele frequency of 0.25. Although the gender-stratified analysis reduced the

statistical power to detect effect of SNPs, we still had 80% of power to detect ORs of 1.43 and 1.44 for men and women respectively. It is important to realise, however, that although the present study involves data on over 3334 individuals, the retrospective and multicentre study design places inevitable limitations on clinical data availability. T2D status and BMI were not available for a substantial subset of MM cases, which did not allow us to adjust our analyses for these variables and, consequently, to rule out the possibility that some of the reported associations could arise as a result of a different distribution of diabetics and/or obese subjects between MM cases and controls. Nonetheless, considering that most of the reported associations with MM risk showed a different direction to those previously published in the GWAS for T2D and given that most of these genes are not linked to obesity, we could not expect to find false positive associations due to these confounding factors.

In conclusion, our study indicate that T2D-related variants within *IGF1*, *KCNJ11*, *CDKN2A-2B*, *MADD*, *THADA*, *LTA*, *FTO*, *ADAM30* and *NOTCH2* genes may influence the risk of MM through insulin-independent mechanisms and that genotyping of specific T2D-related variants may be useful to improve the prediction of MM development. Additional work is needed to replicate our findings in independent and well-characterized populations and functional studies are also warranted to elucidate the biological mechanisms underlying the observed effects.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/ERC-15-0029>.

Declaration of interest

V Andersen is receiving compensation as a consultant for MSD (Merck) and Janssen. The rest of the authors have nothing to disclose.

Funding

This work was supported by grants from the FIBAO foundation (Granada, Spain) and the CRIS foundation against cancer, from the Cancer Network of Excellence (RD12/10 Red de Cáncer) and from the Instituto de Salud Carlos III (Madrid, Spain; PI12/02688).

References

- Alexander DD, Mink PJ, Adami HO, Cole P, Mandel JS, Oken MM & Trichopoulos D 2007 Multiple myeloma: a review of the epidemiologic literature. *International Journal of Cancer* **120** (Suppl 12) 40–61. (doi:10.1002/ijc.22718)
- Altshuler D, Hirschhorn JN, Klannemark M, Lindgren CM, Vohl MC, Nemesh J, Lane CR, Schaffner SF, Bolk S, Brewer C *et al.* 2000 The common PPAR γ Pro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nature Genetics* **26** 76–80. (doi:10.1038/79839)
- Badros A, Golubeva O, Dalal JS, Can I, Thompson J, Rapoport AP, Heyman M, Akpek G & Fenton RG 2007 Neurotoxicity of bortezomib therapy in multiple myeloma: a single-center experience and review of the literature. *Cancer* **110** 1042–1049. (doi:10.1002/cncr.22921)
- Bommert K, Bargou RC & Stuhmer T 2006 Signalling and survival pathways in multiple myeloma. *European Journal of Cancer* **42** 1574–1580. (doi:10.1016/j.ejca.2005.12.026)
- Bouatia-Naji N, Rocheleau G, Van Lommel L, Lemaire K, Schuit F, Cavalcanti-Proença C, Marchand M, Hartikainen AL, Sovio U, De Graeve F *et al.* 2008 A polymorphism within the G6PC2 gene is associated with fasting plasma glucose levels. *Science* **320** 1085–1088. (doi:10.1126/science.1156849)
- Bouatia-Naji N, Bonnefond A, Cavalcanti-Proença C, Sparsø T, Holmkvist J, Marchand M, Delplanque J, Lobbens S, Rocheleau G, Durand E *et al.* 2009 A variant near MTNR1B is associated with increased fasting plasma glucose levels and type 2 diabetes risk. *Nature Genetics* **41** 89–94. (doi:10.1038/ng.277)
- Butt AJ, Firth SM & Baxter RC 1999 The IGF axis and programmed cell death. *Immunology and Cell Biology* **77** 256–262. (doi:10.1046/j.1440-1711.1999.00822.x)
- Cander S, Karkucak M, Gul OO, Sag SO, Yakut T, Ersoy C, Tuncel E & Erturk E 2014 Association between p16(CDKN2A) C540G polymorphism and tumor behavior in prolactinoma: a single-center study. *Biomedical Reports* **2** 589–595. (doi:10.3892/br.2014.281)
- Castillo JJ, Mull N, Reagan JL, Nemr S & Mitri J 2012 Increased incidence of non-Hodgkin lymphoma, leukemia, and myeloma in patients with diabetes mellitus type 2: a meta-analysis of observational studies. *Blood* **119** 4845–4850. (doi:10.1182/blood-2011-06-362830)
- Chambers JC, Elliott P, Zabaneh D, Zhang W, Li Y, Froguel P, Balding D, Scott J & Kooner JS 2008 Common genetic variation near MC4R is associated with waist circumference and insulin resistance. *Nature Genetics* **40** 716–718. (doi:10.1038/ng.156)
- Chen WM, Erdos MR, Jackson AU, Saxena R, Sanna S, Silver KD, Timpson NJ, Hansen T, Orrù M, Grazia Piras M *et al.* 2008 Variations in the G6PC2/ABCB11 genomic region are associated with fasting glucose levels. *Journal of Clinical Investigation* **118** 2620–2628. (doi:10.1172/JCI34566)
- Chen J, Li D, Killary AM, Sen S, Amos CI, Evans DB, Abbruzzese JL & Frazier ML 2009 Polymorphisms of p16, p27, p73, and MDM2 modulate response and survival of pancreatic cancer patients treated with preoperative chemoradiation. *Annals of Surgical Oncology* **16** 431–439. (doi:10.1245/s10434-008-0220-8)
- Chen HY, Huang W, Leung VH, Fung SL, Ma SL, Jiang H & Tang NL 2013 Functional interaction between SNPs and microsatellite in the transcriptional regulation of insulin-like growth factor 1. *Human Mutation* **34** 1289–1297. (doi:10.1002/humu.22363)
- Cheng I, Caberto CP, Lum-Jones A, Seifried A, Wilkens LR, Schumacher FR, Monroe KR, Lim U, Tiirikainen M, Kolonel LN *et al.* 2011 Type 2 diabetes risk variants and colorectal cancer risk: the Multiethnic Cohort and PAGE studies. *Gut* **60** 1703–1711. (doi:10.1136/gut.2011.237727)
- Chiu BC, Gapstur SM, Greenland P, Wang R & Dyer A 2006 Body mass index, abnormal glucose metabolism, and mortality from hematopoietic cancer. *Cancer Epidemiology, Biomarkers & Prevention* **15** 2348–2354. (doi:10.1158/1055-9965.EPI-06-0007)
- Chou YS, Yang CF, Chen HS, Yang SH, Yu YB, Hong YC, Liu CY, Gau JP, Liu JH, Chen PM *et al.* 2012 Pre-existing diabetes mellitus in patients with multiple myeloma. *European Journal of Haematology* **89** 320–327. (doi:10.1111/j.1600-0609.2012.01828.x)
- Colombo M, Mirandola L, Platonova N, Apicella L, Basile A, Figueroa AJ, Cobos E, Chiriva-Internati M & Chiaramonte R 2013 Notch-directed microenvironment reprogramming in myeloma: a single path to multiple outcomes. *Leukemia* **27** 1009–1018. (doi:10.1038/leu.2013.6)

- Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University, and Novartis Institutes of BioMedical Research, Saxena R, Voight BF, Lyssenko V, Burt NP, de Bakker PI, Chen H, Roix JJ, Kathiresan S, Hirschhorn JN *et al.* 2007 Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* **316** 1331–1336. (doi:10.1126/science.1142358)
- Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, Wheeler E, Glazer NL, Bouatia-Naji N, Gloyn AL *et al.* 2010 New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nature Genetics* **42** 105–116. (doi:10.1038/ng.520)
- Duro D, Bernard O, Della Valle V, Berger R & Larsen CJ 1995 A new type of p16INK4/MTS1 gene transcript expressed in B-cell malignancies. *Oncogene* **11** 21–29.
- Feng YH, Velazquez-Torres G, Gully C, Chen J, Lee MH & Yeung SC 2011 The impact of type 2 diabetes and antidiabetic drugs on cancer cell growth. *Journal of Cellular and Molecular Medicine* **15** 825–836. (doi:10.1111/j.1582-4934.2010.01083.x)
- Ferlin M, Noraz N, Hertogh C, Brochier J, Taylor N & Klein B 2000 Insulin-like growth factor induces the survival and proliferation of myeloma cells through an interleukin-6-independent transduction pathway. *British Journal of Haematology* **111** 626–634. (doi:10.1046/j.1365-2141.2000.02364.x)
- Florez JC, Manning AK, Dupuis J, McAteer J, Irenze K, Gianniny L, Mirel DB, Fox CS, Cupples LA & Meigs JB 2007 A 100K genome-wide association scan for diabetes and related traits in the Framingham Heart Study: replication and integration with other genome-wide datasets. *Diabetes* **56** 3063–3074. (doi:10.2337/db07-0451)
- Folsom AR, Pankow JS, Peacock JM, Bielinski SJ, Heiss G & Boerwinkle E 2008 Variation in TCF7L2 and increased risk of colon cancer: the Atherosclerosis Risk in Communities (ARIC) Study. *Diabetes Care* **31** 905–909. (doi:10.2337/dc07-2131)
- Gloyn AL, Weedon MN, Owen KR, Turner MJ, Knight BA, Hitman G, Walker M, Levy JC, Sampson M, Halford S *et al.* 2003 Large-scale association studies of variants in genes encoding the pancreatic β -cell KATP channel subunits Kir6.2 (KCNJ11) and SUR1 (ABCC8) confirm that the KCNJ11 E23K variant is associated with type 2 diabetes. *Diabetes* **52** 568–572. (doi:10.2337/diabetes.52.2.568)
- Grant SF, Thorleifsson G, Reynisdottir I, Benediktsson R, Manolescu A, Sainz J, Helgason A, Stefansson H, Emilsson V, Helgadóttir A *et al.* 2006 Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nature Genetics* **38** 320–323. (doi:10.1038/ng1732)
- Grarup N, Andersen G, Krarup NT, Albrechtsen A, Schmitz O, Jørgensen T, Borch-Johnsen K, Hansen T & Pedersen O 2008 Association testing of novel type 2 diabetes risk alleles in the JAZF1, CDC123/CAMK1D, TSPAN8, THADA, ADAMTS9, and NOTCH2 loci with insulin release, insulin sensitivity, and obesity in a population-based sample of 4,516 glucose-tolerant middle-aged Danes. *Diabetes* **57** 2534–2540. (doi:10.2337/db08-0436)
- Grimberg A 2003 Mechanisms by which IGF-I may promote cancer. *Cancer Biology & Therapy* **2** 630–635. (doi:10.4161/cbt.2.6.678)
- Gudmundsson J, Sulem P, Steinthorsdottir V, Bergthorsson JT, Thorleifsson G, Manolescu A, Rafnar T, Gudbjartsson D, Agnarsson BA, Baker A *et al.* 2007 Two variants on chromosome 17 confer prostate cancer risk, and the one in TCF2 protects against type 2 diabetes. *Nature Genetics* **39** 977–983. (doi:10.1038/ng2062)
- Hamid YH, Urhammer SA, Glümer C, Borch-Johnsen K, Jørgensen T, Hansen T & Pedersen O 2005 The common T60N polymorphism of the lymphotoxin- α gene is associated with type 2 diabetes and other phenotypes of the metabolic syndrome. *Diabetologia* **48** 445–451. (doi:10.1007/s00125-004-1659-1)
- International Myeloma Working Group 2003 Criteria for the classification of monoclonal gammopathies, multiple myeloma and related disorders: a report of the International Myeloma Working Group. *British Journal of Haematology* **121** 749–757. (doi:10.1046/j.1365-2141.2003.04355.x)
- Khan AE, Gallo V, Linseisen J, Kaaks R, Rohrmann S, Raaschou-Nielsen O, Tjønneland A, Johnsen HE, Overvad K, Bergmann MM *et al.* 2008 Diabetes and the risk of non-Hodgkin's lymphoma and multiple myeloma in the European Prospective Investigation into Cancer and Nutrition. *Haematologica* **93** 842–850. (doi:10.3324/haematol.12297)
- Kim WY & Sharpless NE 2006 The regulation of INK4/ARF in cancer and aging. *Cell* **127** 265–275. (doi:10.1016/j.cell.2006.10.003)
- Koh J, Enders GH, Dynlacht BD & Harlow E 1995 Tumour-derived p16 alleles encoding proteins defective in cell-cycle inhibition. *Nature* **375** 506–510. (doi:10.1038/375506a0)
- Kuhn DJ, Berkova Z, Jones RJ, Woessner R, Bjorklund CC, Ma W, Davis RE, Lin P, Wang H, Madden TL *et al.* 2012 Targeting the insulin-like growth factor-1 receptor to overcome bortezomib resistance in preclinical models of multiple myeloma. *Blood* **120** 3260–3270. (doi:10.1182/blood-2011-10-386789)
- Kumar SK, Dispenzieri A, Lacy MQ, Gertz MA, Buadi FK, Pandey S, Kapoor P, Dingli D, Hayman SR, Leung N *et al.* 2014 Continued improvement in survival in multiple myeloma: changes in early mortality and outcomes in older patients. *Leukemia* **28** 1122–1128. (doi:10.1038/leu.2013.313)
- Lau J, Ioannidis JP & Schmid CH 1997 Quantitative synthesis in systematic reviews. *Annals of Internal Medicine* **127** 820–826. (doi:10.7326/0003-4819-127-9-199711010-00008)
- LeRoith D, Werner H, Beitner-Johnson D & Roberts CT Jr 1995 Molecular and cellular aspects of the insulin-like growth factor I receptor. *Endocrine Reviews* **16** 143–163. (doi:10.1210/edrv-16-2-143)
- Li H, Wang J, Mor G & Sklar J 2008 A neoplastic gene fusion mimics trans-splicing of RNAs in normal human cells. *Science* **321** 1357–1361. (doi:10.1126/science.1156725)
- Lope V, Perez-Gomez B, Aragonés N, Lopez-Abente G, Gustavsson P, Plato N, Zock JP & Pollán M 2008 Occupation, exposure to chemicals, sensitizing agents, and risk of multiple myeloma in Sweden. *Cancer Epidemiology, Biomarkers & Prevention* **17** 3123–3127. (doi:10.1158/1055-9965.EPI-08-0343)
- Lyssenko V, Nagorny CL, Erdos MR, Wierup N, Jonsson A, Spégl P, Bugliani M, Saxena R, Fex M, Pulizzi N *et al.* 2009 Common variant in MTNR1B associated with increased risk of type 2 diabetes and impaired early insulin secretion. *Nature Genetics* **41** 82–88. (doi:10.1038/ng.288)
- Ma RC, So WY, Tam CH, Luk AO, Ho JS, Wang Y, Lam VK, Lee HM, Kong AP, Tong PC *et al.* 2014 Genetic variants for type 2 diabetes and new-onset cancer in Chinese with type 2 diabetes. *Diabetes Research and Clinical Practice* **103** 328–337. (doi:10.1016/j.diabres.2013.12.016)
- Mannino GC, Greco A, De Lorenzo C, Andreozzi F, Marini MA, Perticone F & Sesti G 2013 A fasting insulin-raising allele at IGF1 locus is associated with circulating levels of IGF-1 and insulin sensitivity. *PLoS ONE* **8** e85483. (doi:10.1371/journal.pone.0085483)
- Martino A, Sainz J, Buda G, Jamrozik K, Reis RM, Garcia-Sanz R, Jurado M, Ríos R, Szmraj-Rogucka Z, Marques H *et al.* 2012 Genetics and molecular epidemiology of multiple myeloma: the rationale for the IMMENSE consortium (review). *International Journal of Oncology* **40** 625–638. (doi:10.3892/ijo.2011.1284)
- Mohlke KL, Boehnke M & Abecasis GR 2008 Metabolic and cardiovascular traits: an abundance of recently identified common genetic variants. *Human Molecular Genetics* **17** R102–R108. (doi:10.1093/hmg/ddn275)
- Morgan GJ, Walker BA & Davies FE 2012 The genetic architecture of multiple myeloma. *Nature Reviews. Cancer* **12** 335–348. (doi:10.1038/nrc3257)
- Nielsen T, Sparsø T, Grarup N, Jørgensen T, Pisinger C, Witte DR, Diabetes Genetics Replication and Meta-analysis (DIAGRAM) Consortium, Hansen T & Pedersen O 2011 Type 2 diabetes risk allele near CENTD2 is associated with decreased glucose-stimulated insulin release. *Diabetologia* **54** 1052–1056. (doi:10.1007/s00125-011-2054-3)
- Nyholt DR 2004 A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *American Journal of Human Genetics* **74** 765–769. (doi:10.1086/383251)
- Ollberding NJ, Cheng I, Wilkens LR, Henderson BE, Pollak MN, Kolonel LN & Le Marchand L 2012 Genetic variants, prediagnostic circulating levels of insulin-like growth factors, insulin, and glucose and the risk of colorectal cancer: the Multiethnic Cohort study. *Cancer*

- Epidemiology, Biomarkers & Prevention* **21** 810–820. (doi:10.1158/1055-9965.EPI-11-1105)
- Ortega S, Malumbres M & Barbacid M 2002 Cyclin D-dependent kinases, INK4 inhibitors and cancer. *Biochimica et Biophysica Acta* **1602** 73–87.
- Paez D, Gerger A, Zhang W, Yang D, Labonte MJ, Benhanim L, Kahn M, Lenz F, Lenz C, Ning Y et al. 2014 Association of common gene variants in the WNT/ β -catenin pathway with colon cancer recurrence. *Pharmacogenomics Journal* **14** 142–150. (doi:10.1038/tj.2013.20)
- Pan J, Chen C, Jin Y, Fuentes-Mattei E, Velazquez-Tores G, Benito JM, Konopleva M, Andreeff M, Lee MH & Yeung SC 2012 Differential impact of structurally different anti-diabetic drugs on proliferation and chemosensitivity of acute lymphoblastic leukemia cells. *Cell Cycle* **11** 2314–2326. (doi:10.4161/cc.20770)
- Pancewicz J, Taylor JM, Datta A, Baydoun HH, Waldmann TA, Hermine O & Nicot C 2010 Notch signaling contributes to proliferation and tumor formation of human T-cell leukemia virus type 1-associated adult T-cell leukemia. *PNAS* **107** 16619–16624. (doi:10.1073/pnas.1010722107)
- Pechlivanis S, Wagner K, Chang-Claude J, Hoffmeister M, Brenner H & Forsti A 2007 Polymorphisms in the insulin like growth factor 1 and IGF binding protein 3 genes and risk of colorectal cancer. *Cancer Detection and Prevention* **31** 408–416. (doi:10.1016/j.cdp.2007.10.001)
- Prokopenko I, Langenberg C, Florez JC, Saxena R, Soranzo N, Thorleifsson G, Loos RJ, Manning AK, Jackson AU, Aulchenko Y et al. 2009 Variants in MTNR1B influence fasting glucose levels. *Nature Genetics* **41** 77–81. (doi:10.1038/ng.290)
- Qi L, Cornelis MC, Kraft P, Stanya KJ, Linda Kao WH, Pankow JS, Dupuis J, Florez JC, Fox CS, Paré G et al. 2010 Genetic variants at 2q24 are associated with susceptibility to type 2 diabetes. *Human Molecular Genetics* **19** 2706–2715. (doi:10.1093/hmg/ddq156)
- Qian J, Zhou H, Chen J, Ding Q, Cao Q, Qin C, Shao P, Li P, Cai H, Meng X et al. 2014 Genetic polymorphisms in IGF-I and IGFBP-3 are associated with prostate cancer in the Chinese population. *PLoS ONE* **9** e85609. (doi:10.1371/journal.pone.0085609)
- Rattan R, Ali Fehmi R & Munkarah A 2012 Metformin: an emerging new therapeutic option for targeting cancer stem cells and metastasis. *Journal of Oncology* **2012** 928127. (doi:10.1155/2012/928127)
- Richardson PG, Briemberg H, Jagannath S, Wen PY, Barlogie B, Berenson J, Singhal S, Siegel DS, Irwin D, Schuster M et al. 2006 Frequency, characteristics, and reversibility of peripheral neuropathy during treatment of advanced multiple myeloma with bortezomib. *Journal of Clinical Oncology* **24** 3113–3120. (doi:10.1200/JCO.2005.04.7779)
- Richardson PG, Sonneveld P, Schuster MW, Stadtmauer EA, Facon T, Harousseau JL, Ben-Yehuda D, Lonial S, Goldschmidt H, Reece D et al. 2009 Reversibility of symptomatic peripheral neuropathy with bortezomib in the phase III APEX trial in relapsed multiple myeloma: impact of a dose-modification guideline. *British Journal of Haematology* **144** 895–903. (doi:10.1111/j.1365-2141.2008.07573.x)
- Rippe V, Drieschner N, Meiboom M, Murua Escobar H, Bonk U, Belge G & Bullerdiek J 2003 Identification of a gene rearranged by 2p21 aberrations in thyroid adenomas. *Oncogene* **22** 6111–6114. (doi:10.1038/sj.onc.1206867)
- Rung J, Cauchi S, Albrechtsen A, Shen L, Rocheleau G, Cavalcanti-Proença C, Bacot F, Balkau B, Belisle A, Borch-Johnsen K et al. 2009 Genetic variant near IRS1 is associated with type 2 diabetes, insulin resistance and hyperinsulinemia. *Nature Genetics* **41** 1110–1115. (doi:10.1038/ng.443)
- Sainz J, Rudolph A, Hoffmeister M, Frank B, Brenner H, Chang-Claude J, Hemminki K & Forsti A 2012 Effect of type 2 diabetes predisposing genetic variants on colorectal cancer risk. *Journal of Clinical Endocrinology and Metabolism* **97** E845–E851. (doi:10.1210/jc.2011-2565)
- Sandhu MS, Weedon MN, Fawcett KA, Wasson J, Debenham SL, Daly A, Lango H, Frayling TM, Neumann RJ, Sherva R et al. 2007 Common variants in WFS1 confer risk of type 2 diabetes. *Nature Genetics* **39** 951–953. (doi:10.1038/ng2067)
- Sauroja I, Smeds J, Vlaykova T, Kumar R, Talve L, Hahka-Kemppinen M, Punnonen K, Jansen CT, Hemminki K & Pyrhonen S 2000 Analysis of G(1)/S checkpoint regulators in metastatic melanoma. *Genes, Chromosomes & Cancer* **28** 404–414. (doi:10.1002/1098-2264(200008)28:4<404::AID-GCC6>3.0.CO;2-P)
- Saxena R, Hivert MF, Langenberg C, Tanaka T, Pankow JS, Vollenweider P, Lyssenko V, Bouatia-Naji N, Dupuis J, Jackson AU et al. 2010 Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nature Genetics* **42** 142–148. (doi:10.1038/ng.521)
- Scott LJ, Bonnycastle LL, Willer CJ, Sprau AG, Jackson AU, Narisu N, Duren WL, Chines PS, Stringham HM, Erdos MR et al. 2006 Association of transcription factor 7-like 2 (TCF7L2) variants with type 2 diabetes in a Finnish sample. *Diabetes* **55** 2649–2653. (doi:10.2337/db06-0341)
- Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, Erdos MR, Stringham HM, Chines PS, Jackson AU et al. 2007 A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* **316** 1341–1345. (doi:10.1126/science.11442382)
- Scuteri A, Sanna S, Chen WM, Uda M, Albai G, Strait J, Najjar S, Nagaraja R, Orru M, Usala G et al. 2007 Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLoS Genetics* **3** e115. (doi:10.1371/journal.pgen.0030115)
- Sesti G, Mannino GC, Andreozzi F, Greco A, Perticone M, Sciacqua A, Marini MA & Perticone F 2014 A polymorphism at IGF1 locus is associated with carotid intima media thickness and endothelium-dependent vasodilatation. *Atherosclerosis* **232** 25–30. (doi:10.1016/j.atherosclerosis.2013.10.024)
- Sharifi S, Daghighi S, Motazacker MM, Badlou B, Sanjabi B, Akbarkhanzadeh A, Rowshani AT, Laurent S, Peppelenbosch MP & Rezaee F 2013 Superparamagnetic iron oxide nanoparticles alter expression of obesity and T2D-associated risk genes in human adipocytes. *Scientific Reports* **3** 2173. (doi:10.1038/srep02173)
- Shu XO, Long J, Cai Q, Qi L, Xiang YB, Cho YS, Tai ES, Li X, Lin X, Chow WH et al. 2010 Identification of new genetic risk variants for type 2 diabetes. *PLoS Genetics* **6** e1001127. (doi:10.1371/journal.pgen.1001127)
- Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin P, Vincent D, Belisle A, Hadjadj S et al. 2007 A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* **445** 881–885. (doi:10.1038/nature05616)
- Sprynski AC, Hose D, Caillot L, Reme T, Shaughnessy JD Jr, Barlogie B, Seckinger A, Moreaux J, Hundemer M, Jourdan M et al. 2009 The role of IGF-1 as a major growth factor for myeloma cell lines and the prognostic relevance of the expression of its receptor. *Blood* **113** 4614–4626. (doi:10.1182/blood-2008-07-170464)
- Standal T, Borset M, Lenhoff S, Wisloff F, Stordal B, Sundan A, Waage A & Seidel C 2002 Serum insulinlike growth factor is not elevated in patients with multiple myeloma but is still a prognostic factor. *Blood* **100** 3925–3929. (doi:10.1182/blood-2002-05-1406)
- Steinthorsdottir V, Thorleifsson G, Reynisdottir I, Benediktsson R, Jonsdottir T, Walters GB, Stykarsdottir U, Gretarsdottir S, Emilsson V, Ghosh S et al. 2007 A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. *Nature Genetics* **39** 770–775. (doi:10.1038/ng2043)
- Stevens VL, Ahn J, Sun J, Jacobs EJ, Moore SC, Patel AV, Berndt SI, Albanes D & Hayes RB 2010 HNF1B and JAZF1 genes, diabetes, and prostate cancer risk. *Prostate* **70** 601–607. (doi:10.1002/pros.21094)
- Takeuchi F, Serizawa M, Yamamoto K, Fujisawa T, Nakashima E, Ohnaka K, Ikegami H, Sugiyama T, Katsuya T, Miyagishi M et al. 2009 Confirmation of multiple risk loci and genetic impacts by a genome-wide association study of type 2 diabetes in the Japanese population. *Diabetes* **58** 1690–1699. (doi:10.2337/db08-1494)
- Tang Y, Han X, Sun X, Lv C, Zhang X, Guo W, Ren Q, Luo Y, Zhang X, Zhou X et al. 2013 Association study of a common variant near IRS1 with type 2 diabetes mellitus in Chinese Han population. *Endocrine* **43** 84–91. (doi:10.1007/s12020-012-9693-0)
- Than BL, Goos JA, Sarver AL, O'Sullivan MG, Rod A, Starr TK, Fijneman RJ, Meijer GA, Zhao L, Zhang Y et al. 2013 The role of KCNQ1 in mouse and

- human gastrointestinal cancers. *Oncogene* **33** 3861–3868. (doi:10.1038/onc.2013.350)
- Tsai FJ, Yang CF, Chen CC, Chuang LM, Lu CH, Chang CT, Wang TY, Chen RH, Shiu CF, Liu YM *et al.* 2010 A genome-wide association study identifies susceptibility variants for type 2 diabetes in Han Chinese. *PLoS Genetics* **6** e1000847. (doi:10.1371/journal.pgen.1000847)
- Turner A, Li LC, Pilli T, Qian L, Wiley EL, Setty S, Christov K, Ganesh L, Maker AV, Li P *et al.* 2013 MADD knock-down enhances doxorubicin and TRAIL induced apoptosis in breast cancer cells. *PLoS ONE* **8** e56817. (doi:10.1371/journal.pone.0056817)
- Unoki H, Takahashi A, Kawaguchi T, Hara K, Horikoshi M, Andersen G, Ng DP, Holmkvist J, Borch-Johnsen K, Jørgensen T *et al.* 2008 SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nature Genetics* **40** 1098–1102. (doi:10.1038/ng.208)
- Voight BF, Scott LJ, Steinthorsdottir V, Morris AP, Dina C, Welch RP, Zeggini E, Huth C, Aulchenko YS, Thorleifsson G *et al.* 2010 Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nature Genetics* **42** 579–589. (doi:10.1038/ng.609)
- Wallin A & Larsson SC 2011 Body mass index and risk of multiple myeloma: a meta-analysis of prospective studies. *European Journal of Cancer* **47** 1606–1615. (doi:10.1016/j.ejca.2011.01.020)
- Watson J 2013 Oxidants, antioxidants and the current incurability of metastatic cancers. *Open Biology* **3** 120144. (doi:10.1098/rsob.120144)
- Wellcome Trust Case Control C 2007 Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* **447** 661–678. (doi:10.1038/nature05911)
- Willer CJ, Bonnycastle LL, Conneely KN, Duren WL, Jackson AU, Scott LJ, Narisu N, Chines PS, Skol A, Stringham HM *et al.* 2007 Screening of 134 single nucleotide polymorphisms (SNPs) previously associated with type 2 diabetes replicates association with 12 SNPs in nine genes. *Diabetes* **56** 256–264. (doi:10.2337/db06-0461)
- Wu W, Merriman K, Nabaah A, Seval N, Seval D, Lin H, Wang M, Qazilbash MH, Baladandayuthapani V, Berry D *et al.* 2014 The association of diabetes and anti-diabetic medications with clinical outcomes in multiple myeloma. *British Journal of Cancer* **111** 628–636. (doi:10.1038/bjc.2014.307)
- Xu F, Gardner A, Tu Y, Michl P, Prager D & Lichtenstein A 1997 Multiple myeloma cells are protected against dexamethasone-induced apoptosis by insulin-like growth factors. *British Journal of Haematology* **97** 429–440. (doi:10.1046/j.1365-2141.1997.592708.x)
- Xu D, Hu J, Xu S, De Bruyne E, Menu E, Van Camp B, Vanderkerken K & Van Valckenborgh E 2012a Dll1/Notch activation accelerates multiple myeloma disease development by promoting CD138+ MM-cell proliferation. *Leukemia* **26** 1402–1405. (doi:10.1038/leu.2011.332)
- Xu D, Hu J, De Bruyne E, Menu E, Schots R, Vanderkerken K & Van Valckenborgh E 2012b Dll1/Notch activation contributes to bortezomib resistance by upregulating CYP1A1 in multiple myeloma. *Biochemical and Biophysical Research Communications* **428** 518–524. (doi:10.1016/j.bbrc.2012.10.071)
- Xu CX, Zhu HH & Zhu YM 2014 Diabetes and cancer: associations, mechanisms, and implications for medical practice. *World Journal of Diabetes* **5** 372–380. (doi:10.4239/wjd.v5.i3.372)
- Yamauchi T, Hara K, Maeda S, Yasuda K, Takahashi A, Horikoshi M, Nakamura M, Fujita H, Grarup N, Cauchi S *et al.* 2010 A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B. *Nature Genetics* **42** 864–868. (doi:10.1038/ng.660)
- Yasuda K, Miyake K, Horikawa Y, Hara K, Osawa H, Furuta H, Hirota Y, Mori H, Jonsson A, Sato Y *et al.* 2008 Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nature Genetics* **40** 1092–1097. (doi:10.1038/ng.207)
- Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H, Timpson NJ, Perry JR, Rayner NW, Freathy RM *et al.* 2007 Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science* **316** 1336–1341. (doi:10.1126/science.1142364)
- Zeggini E, Scott LJ, Saxena R, Voight BF, Marchini JL, Hu T, de Bakker PI, Abecasis GR, Almgren P, Andersen G *et al.* 2008 Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nature Genetics* **40** 638–645. (doi:10.1038/ng.120)

Received in final form 7 May 2015

Accepted 15 May 2015